



PACS – A Photodetector Array Camera & Spectrometer for Herschel



Instrument Design

Projection of focal plane onto spectrometer arrays



Integral Field Spectrometer


Simultaneous 57-105 & 105-210 μm spectroscopy.

47" x 47" (5x5 pixels) FOV rearranged via an image slicer on two 16x25 Ge:Ga detector arrays.



$$\lambda/\Delta\lambda \sim 1500$$

Sensitivity: $\sim 5 \times 10^{-18} \text{ W/m}^2$ (5σ , 1h)

PACS is one of three science instruments for ESA's Herschel mission. It operates either as an imaging photometer or an integral field spectrometer over the spectral band from 57 to 210 μm .



Projection of focal plane onto bolometer arrays



Red photometer
32x16 pixels (2x16x16)

Blue photometer
64x32 pixels (4x2x16x16)

Imaging Photometer

Simultaneous two-band (same FOV) 60-85 μm or 85-130 μm and 130-210 μm imaging.

Two filled bolometer arrays: 32x16 and 64x32 pixels


Point source detection limit: $\sim 3 \text{ mJy}$ (5σ , 1h)




PACS is being designed and built by a consortium of institutes and university departments from across Europe under the leadership of Principal Investigator Albrecht Poglitsch located at Max-Planck-Institute for Extraterrestrial Physics, Garching, Germany. Consortium members are: Austria: UVIE; Belgium: IMEC, KUL, CSL; France: CEA, OAMP; Germany: MPE, MPIA; Italy: IFSI, OAP/OAT, OAA/CAISMI, LENS, SISSA; Spain: IAC.



For more information please visit: <http://pacs.mpe.mpg.de/> or <http://www.rssd.esa.int/> or <http://www.ipac.caltech.edu/>



PACS – A Photodetector Array Camera & Spectrometer for Herschel



Scientific Motivations

The opening of the 60-210 μm window by PACS to sensitive photometry and spectroscopy at high spatial resolution will address a wide range of key questions of current astrophysics concerning the ***origins of stars, planetary systems, galaxies, and the evolution of the Universe.***


-Most of the energy released e.g. in starbursts or AGNs is absorbed by interstellar dust (which prevents observation at shorter wavelengths) and re-emitted in the far infrared and sub-mm domain.

-Cool, dusty and/or distant objects have their emission peak in the far-IR.

-The far-IR also contains many spectral lines from atoms, ions and molecules. Largely unaffected by extinction they provide detailed information on UV radiation, density, temperature, velocities and abundances of ionised and neutral components of interstellar and circumstellar gas.

Some examples:

What is the cosmic history of star formation and AGN activity?



Simulated deep PACS survey of 10^{-5} sr at 75, 110 and 175 μm (false colors) to a 1σ limit of ~ 0.5 mJy (50h). Deepest sources are at $z\sim 3$.

- Deep multi-band photometric surveys and spectroscopy at the peak of cosmic star formation ($z=1\dots 3$)


How does stellar mass loss influence the ISM chemistry?

The carbon star YCVn. ISOPHOT 90 μm map (Izumiura et al. 1996)




- Photometric mapping and spectroscopy (e.g. CO, H₂O, OI) of the circumstellar matter in evolved objects

How do stars form out of the interstellar medium?



M82 (Subaru/FOCAS) with the PACS spectroscopy FOV overlaid

- Local galaxies: photometric and spectral line mapping for detailed, spatially resolved studies of star formation on galactic scales



ρ Oph, SCUBA 850 μm (Johnstone et al. 2000)

- Photometric surveys of nearby molecular clouds: search for protostars

PACS is also intended to be an important driver for other projects which will explore adjacent spectral regions, such as JWST in the near/mid IR and ALMA in the mm domain.