

TNOs are Cool: A Survey of the Transneptunian Region

Principal Investigator: Thomas Mueller

Institution: MPE, Garching

E-mail address: tmueller@mpe.mpg.de

Co-Investigators: Emmanuel Lellouch (Co-PI), Observ. de Paris-Meudon
Hermann Bönhardt (Co-PI), MPS, Katlenburg-Lindau
Antonella Barucci, Observatoire de Paris-Meudon
Jacques Crovisier, Observatoire de Paris-Meudon
Audrey Delsanti, Observatoire de Paris-Meudon
Alain Doressoundiram, Observatoire de Paris-Meudon
Elisabetta Dotto, Osservatorio Astronomico di Roma
Sonia Fornasier, Observatoire de Paris-Meudon
Olivier Groussin, Lab. d'Astrophysique de Marseille
Pedro J. Gutiérrez, IAA-CSIC, Granada
Olivier Hainaut, ESO, Chile
Alan Harris, DLR, Berlin
Paul Hartogh, MPS, Katlenburg-Lindau
Daniel Hestroffer, Observatoire de Paris
Jonathan Horner, Open University, Milton Keynes
Dave Jewitt, University of Hawaii
Mark Kidger, ESAC, Villafranca del Castillo
Csaba Kiss, Konkoly Observatory, Budapest
Pedro Lacerda, University of Hawaii
Luisa Lara, IAA-CSIC, Granada
Tanya Lim, RAL, Didcot
Raphael Moreno, Observatoire de Paris-Meudon
Michael Mueller, University of Arizona
José Luis Ortiz, IAA-CSIC, Granada
Miriam Rengel, MPS, Katlenburg-Lindau

Pablo Santos–Sanz, IAA–CSIC, Granada

John Stansberry (NASA PI), University of Arizona

Bruce Swinyard, RAL, Didcot

Nicolas Thomas, University of Bern

David Trilling, University of Arizona

Science Category: Solar System: Kuiper Belt bodies

Observing Modes: Pacs Photometer, Spire Photometer

Hours Requested: 378.5

Abstract:

Over one thousand objects have been discovered orbiting beyond Neptune. These trans–Neptunian objects (TNOs) represent the primitive remnants of the planetesimal disk from which the outer planets formed, and is an analog for unseen dust parent–bodies in debris disks observed around other main–sequence stars. The dynamical and physical properties of these bodies provide unique and important constraints on formation and evolution models of the outer Solar System. While the dynamical architecture in this region (also known as the Kuiper Belt) is becoming relatively clear, the physical properties of the objects are only beginning to be revealed. In particular, fundamental parameters such as size, albedo, density and thermal properties are difficult to measure. Measurements of their thermal emission, which peaks at far–IR wavelengths, offer the best means available to determine those physical properties. While Spitzer has provided the first results, notably revealing a large albedo diversity in this population, the increased sensitivity of Herschel and its wavelength coverage will permit profound advances in the field. We propose to perform radiometric measurements of 141 objects, including 25 known multiple systems. This large sample will permit: (i) A determination of the size distribution of the large (> 200 km) objects, thought to have remained unchanged from the accretion phase. (ii) Systematic searches for correlations between size, albedo, and other physical and orbital parameters, diagnostic of formation and evolution processes. (iii) Determination of mass–density for at least 20 binary TNOs, diagnostic of nebular chemistry and interior structure. (iv) The first study of their thermophysical properties, including thermal inertia and surface emissivity. When combined with measurements of the dust population beyond Neptune (e.g. from the New Horizons mission to Pluto and beyond), our results will provide a benchmark for understanding the solar system debris disk, and extra–solar ones as well.