

## D Exercises 4

### D.1 The mass of Sgr A\*

Derive the formula relating the mass of the central object with semi-major axis and orbital period. Then plug in some values for Sgr A\*:  $P = 16.0$  yr,  $a = 125$  mas (milli-arcsec), and distance  $R_0 = 8.3$  kpc. You will also need the gravitational constant  $G = 6.67 \times 10^{-11} \text{ N m}^2/\text{kg}^2$ . (Express the result in solar masses,  $M_\odot = 2 \times 10^{30} \text{ kg}$ .)

### D.2 Density

What is the density of a black hole as a function of mass, if one identifies its size with the Schwarzschild radius? At what mass is the density of a black hole smaller than that of air? (Express the result in solar masses,  $M_\odot = 2 \times 10^{30} \text{ kg}$ .)

### D.3 Gaussian signals

Show that for a Gaussian signal

$$P_G(x) = \frac{1}{2\pi\sqrt{\sigma}} \exp\left(-\frac{x^2}{2\sigma^2}\right)$$

The following equality holds:

$$\begin{aligned} \langle \exp(\alpha x) \rangle &:= \int \exp(\alpha x) P_G(x) dx \\ &= \exp\left(\frac{1}{2}\alpha^2 \langle x^2 \rangle\right) \end{aligned}$$

### D.4 Structure function and coherence function

Show that for a real-valued function  $A$  the structure function  $D_A$  and coherence function  $B_A$

$$\begin{aligned} D_A(\vec{r}) = D_A(\vec{r}, \vec{0}) &= \langle |A(\vec{r}) - A(\vec{0})|^2 \rangle \\ B_A(\vec{r}) &= \langle A(\vec{r}) A^*(\vec{r}) \rangle \end{aligned}$$

have the simple relation

$$D_A(\vec{r}) = 2(B_A(\vec{0}) - B_A(\vec{r}))$$

Here

$$\langle Y(\vec{x}) \rangle = \int Y(\vec{x}, \vec{r}) d^3r$$

(This relation is used twice in the derivation of the Fried parameter.)

## D.5 Radial free-fall in Newtonian gravity

Solve the equation for radial infall in Newtonian gravity to get  $t(r)$ .

- Start from the force equation
- Transform from  $\ddot{r}$  to  $dv/dr$ .
- Solve the (very simple) differential equation to get  $v(r)$ .
- Fix the integration constant by demanding that at the start point  $r_B$ ,  $v(r_B) = 0$ .
- Now use  $v = dr/dt$  and separate variables to get an expression for  $t(r)$ .
- The integral is some work - either look it up, or try with substitutions...
- When does that particle reach  $r = 0$  (as a function of  $r_B$  and  $M$ )
- Starting from  $r_B = 1 \text{ AU}$  for  $M = 1 M_\odot$ , when does the test particle reach the center?
- The motion can also be seen as an  $e = 1$  orbit with a semi-major axis of  $a = r_B/2$ . The time to reach the center is thus half of the the orbital period for that orbit. Does that agree with what you calculated before? It should.