

# Thermal Emission from Isolated Neutron Stars

- Theoretical aspects
- Observations and model applications

Slava Zavlin (MPE, Garching)




# A Short History

- Chi & Salpeter (1964) and Tsuruta (1964):  
thermal radiation from the surface of a hot NS  
may be a source of cosmic X-rays
- First detections with Einstein (1978-81) and EXOSAT (1983-86):  
middle-aged PSRs B0656+14 and B1055-52  
old PSRs B0950+08 and B1929+10  
central compact sources in the SNRs RCW 103, Puppis A, Kes 73 and  
PKS 1209-52
- X-ray studies in the “Decade of Space Science” since 1991 with  
ROSAT, ASCA, EUVE, BeppoSax, Chandra and XMM-Newton

In total thermal emission detected from about a dozen of isolated NSs:

- from the whole surface - atmosphere - of cooling NSs
- from polar caps heated by relativistic particles streaming down into  
the surface from pulsar's magnetosphere

# Thermal vs. Nonthermal Emission in Pulsars of Different Ages



**Young** active ( $<1$  kyr):  
**nonthermal** radiation  
dominates

**Middle-aged** ( $10-10^2$  kyr):  
**thermal** component in soft  
EUV and soft X-rays

**Old** ( $>10^3$  kyr):  
**thermal** emission in  
optical-UV and/or  
soft X-rays

- No **nonthermal** radiation is seen in "dead" pulsars

# First Questions

- Why studying thermal emission is needed?
- Why not to take the blackbody model?
- What is the state of the NS surface?  
Isn't it just solid?
- What is the chemical composition of the NS surface?  
Is it hydrogen?  
Or it's composed of heavier elements?

## ● Why studying thermal emission is needed?

Comparing observed emission with theoretical models  $\Rightarrow$

$T_{\text{sur}}, B, R, M$

$T_{\text{sur}} \Rightarrow$  thermal evolution

$R, M \Rightarrow$  constraints on **EOS** and internal structure

surface **chemical** composition  $\Rightarrow$

**formation** of NSs and their **interaction** with environment

## ● Why not to take the blackbody model?

Stars are **not** black bodies

Thermal radiation  $\neq$  blackbody radiation

● What is the state of the NS surface? Isn't it just solid?

It depends on  $T_{\text{sur}}$ ,  $B$  and chemical composition.

For hydrogen, the surface is in a condensed state if:

$$\begin{aligned} T_{\text{sur}} < 1 \times 10^5 \text{ K} & \quad \text{at} \quad B = 1 \times 10^{13} \text{ G} \\ T_{\text{sur}} < 5 \times 10^5 \text{ K} & \quad \text{at} \quad B = 1 \times 10^{14} \text{ G} \\ T_{\text{sur}} < 1 \times 10^6 \text{ K} & \quad \text{at} \quad B = 5 \times 10^{14} \text{ G} \quad (\text{Lai \& Salpeter 1997}) \end{aligned}$$

● What is the **chemical** composition of the NS surface?

Is it **hydrogen**? Or it's composed of heavier elements?

A small amount of H,  $\rho \approx 10^{-3} - 10^{-1} \text{ g/cm}^2$  due to accretion from ISM or fallback of material ejected during the SNR explosion, determines properties of the atmosphere.

Nuclear spallation reactions may destroy heavy elements (Bildsten et al. 1992, 1993).

Othewise, **other** elements may be present.

# Main Aspects of the NS Atmosphere Modeling

● What's special about NS atmospheres?

Why not to use standart stellar atmosphere models?

1. **Enormous gravity** at the surface ( $M \approx 1.4 M_{\odot}$ ,  $R \approx 10$  km)

$$g \approx 10^{14} \text{ vs. } 10^4 \text{ cm}^2/\text{s} \text{ for usual stars}$$

⇒ NS atmospheres are strongly **compressed**

$$\rho \approx 10^{-2} - 10^1 \text{ vs. } 10^{-7} \text{ g/cm}^3$$

$$H \approx kT_{\text{sur}}/m_{\text{p}}g \approx 10^{-1} - 10^1 \text{ vs. } 10^8 \text{ cm}$$

⇒ **stratification** of chemical elements

⇒ **nonideality** effects (pressure ionization, smoothed spectral features)

## 2. Huge magnetic fields, $B \approx 10^{10} - 10^{14} \text{ G}$

$\Rightarrow E_{CE} = 11.6 (B/10^{12} \text{ G}) \text{ keV} \gg kT_{\text{sur}} \approx 0.1 - 1 \text{ keV}$

$\Rightarrow$  NS atmospheres are essentially **anisotropic**

$\Rightarrow$  opacities depend on the direction and polarisation of radiation

$\Rightarrow$  radiation is **polarized**

$\Rightarrow \gamma = E_{CE}/(Z^2 \text{ Ry}) = 850 Z^{-2} (B/10^{12} \text{ G}) \gg 1$

$\Rightarrow$  **atomic structure** is distorted by  $B$

$\Rightarrow$  increase of binding (ionization) energies of bound states

$$I/(Z^2 \text{ Ry}) \approx \ln^2(\gamma/Z^2) \gg 1, \quad I_H \approx 0.2 \text{ keV at } B = 10^{13} \text{ G}$$

$\Rightarrow$  altered ionisation equilibrium

$\Rightarrow$  radiation depends on  $B$


$\Rightarrow$  proton cyclotron line at  $E_{CP} = 63 (B/10^{14} \text{ G}) \text{ eV}$




3. Nonuniform magnetic field  $\Rightarrow$   
nonuniform surface temperature distribution  $\Rightarrow$   
thermal radiation is pulsed
  
4. Gravitational bending of photon trajectories strongly affects the shape of pulsations

# Gravitational bending of photon trajectories



$$g_r = (1 - R_G/R)^{1/2}, \quad R_G = 2GM/c^2$$




the whole surface is visible if  $g_r < 0.69$




# NS atmosphere models with low magnetic fields, $B < 10^8 - 10^9 \text{ G}$






- radiation is essentially anisotropic
- anisotropy depends on energy

# Spectra of nonmagnetic NS atmospheres with various abundances of heavy elements



# Effect of the surface gravity $g$



larger  $g \Rightarrow$



higher density  $\rho \Rightarrow$


stronger ionization  $\Rightarrow$

*weaker* features

(important at low  $T_{eff}$ )

# NS atmosphere models with strong magnetic fields, $B > 10^{11}$ G






Angular dependence of emitted radiation:

"pencil"-like structure along B

"fan"-like structure across B





**proton cyclotron line**  
 $B = 10^{14} \text{ G}$




**atomic transitions**  
 $T_{\text{eff}} < 10^6 \text{ K}$


**electron cyclotron line**  
 $B = 3 \times 10^{11} \text{ G}$



# Magnetic iron spectra (Rajagopal et al. 1997)




Emission from the **whole** NS surface:  
**smearing** of spectral features due to fast rotation




$P < 10$  ms

# Light curves of radiation from a magnetized NS



# NS atmosphere vs. blackbody model



$$T_{\text{bb}} > T_{\text{atm}}$$

$$S_{\text{bb}} < S_{\text{atm}}$$

# Atmosphere Models vs. Multiwavelength Observational Data on Isolated Neutron Stars

- the millisecond pulsar [J0437-4715](#)
- the [Vela](#) pulsar
- the middle-aged pulsars [B0656+14](#) and [B1055-52](#)
- the X-ray pulsar [1E 1207.4-5209](#) in the SNR [PKS 1209-51](#)
- some others...

ROSAT and Chandra on  
ms-pulsar J0457-4715:


thermal emission from PCs with  
nonuniform temperature  $\Rightarrow$   
bulk of X-rays

plus


nonthermal PL ( $E > 2$  keV)

or

a single broken PL  
with  $E_{br} \approx 1.7$  keV ?



# HST data on PSR J0437-4715 in far-UV



support:

X-ray PCs+PL plus emission from the rest surface of  $T_{\text{sur}} \approx 6 \times 10^4$  K

rule out:


the broken PL model

PSR J003+0451:

optical data rule out the nonthermal interpretation of the X-ray emission (Koptsevich et al. 2003)



# Chandra Resolves the *Vela* Pulsar from Its Nebula



# The Vela pulsar

LETGS: a smooth thermal spectrum at  $E < 2$  keV

ACIS: second nonthermal component

Thermal component:

blackbody model —

$R_{bb} \approx 2.5$  km at  $d \approx 300$  pc (parallax)

$T_{bb} \approx 3 \times 10^6$  K

with PL of  $\gamma \approx 2.8$


or

hydrogen magnetized NS atmosphere —


$d \approx 250$  pc at  $R = 10$  km

$T \approx 9 \times 10^5$  K

with PL of  $\gamma \approx 1.5$



The multiwavelength spectrum of the *Vela* pulsar:  
interpretation involving the NS atmosphere model



More (successful) **NS atmosphere** model applications:

- radio-silent NS **RX J0822-4300** in the SNR Puppis A  
⇒ distance consistent with independent measurements  
(blackbody model ⇒ 6 times larger distance)


- transiently accreting NSs in X-ray binaries

**Aql X-1, Cen X-4, KS 1713-260, 4U 2129+47**

**quiescent** radiation from NS hydrogen atmospheres

(Rutledge et al. 1999-2002; Nowak et al. 2002)

due to heat released in the compressed material



Middle-aged PSR B0656+14:

two thermal components,  
blackbody model:

"soft"

$$T_{\text{bb}} \approx 8.5 \times 10^5$$

$$R_{\text{bb}} \approx 18$$

"hard"

$$1.7 \times 10^6 \text{ K}$$

$$1.5 \text{ km}$$

$$(d \approx 750 \text{ pc})$$


plus a PL of  $\gamma \approx 1.5$

NS atmosphere models:

too small distance,  $d \approx 100 \text{ pc}$   
for  $R=10 \text{ km}$

Chandra LETGS and ACIS data

# Multiwavelength spectrum of PSR B0656+14



## Middle-aged PSR B1055-52

Similar three-component interpretation, TS+TH+PL:

PL dominates at  $E > 2$  keV, fits also the optical and  $\gamma$ -ray fluxes

PSRs	B0656+14	B1055-52
"soft" $T_{\text{bb}}$	$8.5 \times 10^5$ K	$7.5 \times 10^5$ K
age $\tau = P / (2dP/dt)$	110 kyr	540 kyr


- the mass of B1055-52 is smaller (Yakovlev et al. 2002)  $\Rightarrow$  lower central density and weaker neutrino emission  $\Rightarrow$  slower cooling
- characteristic ages are **not** true ages

# Light curves of X-rays from PSR B0656+14:

strong energy dependence of pulse shape and pulsed fraction  $p_f$

$p_f$  ↓ from 18% at 0.2 keV to 9% at 0.6 keV

⇒ blackbody model does **not** work (Page 1995)





## Problems:

- middle-aged PSRs **B0656+14** and **B1055-52** —

**NS atmosphere models** are **not** applicable  $\Rightarrow$

**no** atmospheres ?

too cold and/or too strong magnetic field?

blackbody model works on spectral data but

is **not** able to explain the temporal behavior

(radiation has to be **anisotropic**)


- “truly” isolated, “dim” radio-quiet NSs (old and rather cold)

**RX J1856-3754**, **J0720-3125** (and a few more)

**none** of the standard NS atmosphere works  $\Rightarrow$

**condensed**-matter surface? or **thin** atmosphere?

# 1E 1207.4-5209 in the SNR PKS 1209-51: ROSAT (1993)



Chandra:

smooth pulsations with

$$p_f \approx 6\%$$

$$P = 0.42 \text{ s}$$

$$dP/dt \approx 2 \times 10^{-13} \text{ s s}^{-1}$$

confirmed with XMM


$$\Rightarrow B \approx 3 \times 10^{12} \text{ G}$$

$$\tau \approx 200\text{-}1600 \text{ kyr}$$

vs.

7-20 kyr for the SNR

# Chandra X-ray spectrum of 1E 1207.4-5209



Two broad absorption lines at 0.7 and 1.4 keV —  
first direct evidence of a NS atmosphere:


no hydrogen

once-ionized helium at  $B \approx 10^{14}$  G?

He-like oxygen or neon at  $B \leq 10^{12}$  G (Mori & Hailey 2002)?

new XMM (260 ks, done) and Chandra (300 ks, scheduled) observations

# Spectra of 1E 1207.4-5209 at two rotational phases:



the lines are stronger at  
the pulse minimum

## (Some) Conclusions

- More data  $\Rightarrow$  more questions:

true ages of NSs?

surface temperature and magnetic field distributions?

actual mechanism(s) of the surface emission?

states of the surface matter?

.....

- Any nonstandard approaches are welcome

E. g., plate tectonics (Ruderman 1991) —

magnetized and hot plates on the nonmagnetic and cold surface

(Page et al. 1995)

or.....your suggestions?